

(\*) **Definitions**

1. The function  $F(x)$  is an **antiderivative** of the function  $f(x)$  if  $F'(x) = f(x)$ .
2. The collection of all antiderivatives of a function  $f(x)$  is called the **indefinite integral** of  $f(x)$ ,<sup>†</sup> and denoted by

$$\int f(x) dx.$$

(\*) **Comments/Facts**

- (i) Every continuous function has antiderivatives.
- (ii) If a function  $f(x)$  has the antiderivative  $F(x)$ , then for any constant  $C$ , the function  $F(x) + C$  is also an antiderivative of  $f(x)$ . Quasi-conversely, if  $F(x)$  and  $G(x)$  are both antiderivatives of  $f(x)$ , then  $G(x) = F(x) + C$  for some constant  $C$ . In other words, if  $f(x)$  has an antiderivative, then it has infinitely many and they all have the form  $F(x) + C$ , where  $F(x)$  is any one antiderivative of  $f(x)$ .
- (iii) In terms of the notation for the indefinite integral of  $f(x)$ , we write

$$\int f(x) dx = F(x) + C,$$

where  $F(x)$  is any one antiderivative of  $f(x)$  and  $C$  (the *constant of integration*) is an unspecified constant that can take any value.

- (iv) The reason that a factor of  $dx$  is included in the indefinite integral is due to the (deep and important) relation between antiderivatives and **definite integrals**, about which we will learn in a week or so.

(\*) **Integration Rules/Formulas** (the basic ones)

$$(a) \int f(x) \pm g(x) dx = \left( \int f(x) dx \right) \pm \left( \int g(x) dx \right),$$

i.e., *the integral of a sum is equal to the sum of the integrals.*

$$(b) \int a f(x) dx = a \int f(x) dx,$$

i.e., *the integral of a constant multiple is the constant multiple of the integral, where  $a$  is a **nonzero** constant.*

$$(c) \int x^k dx = \frac{x^{k+1}}{k+1} + C, \text{ for all } k \neq -1. \text{ The case } k = -1 \text{ is covered next.}$$

$$(d) \int x^{-1} dx = \int \frac{dx}{x} = \ln|x| + C.$$

---

<sup>†</sup>Technically it is the indefinite integral of the differential  $f(x) dx$ .